

Explicit estimators under m -dependence for a multivariate normal distribution

Martin Ohlson¹, Zhanna Andrushchenko²,
and Dietrich von Rosen²

¹Linköping University, Sweden

²Swedish University of Agricultural Sciences, Uppsala, Sweden

Abstract

The problem of estimating parameters of a multivariate normal p -dimensional random vector is considered for a banded covariance structure under m -dependence. A simple non-iterative estimation procedure is suggested which gives explicit, unbiased and consistent estimator of the mean and explicit and consistent estimator of the covariance matrix for arbitrary p and m .

Furthermore, this estimation procedure and a discussion about the residuals are used for estimation of the mean and the covariance matrix in a Growth Curve model again when the covariance matrix is banded of order m .

Keywords

Multivariate normal distribution, Banded covariance matrices, Covariance matrix estimation, Explicit estimators, Growth curve model, Residuals.

References

- Ohlson, M., Z. Andrushchenko, and D. von Rosen (2007). Explicit estimators under m -dependence for a multivariate normal distribution. Submitted to *Ann. Inst. Statist. Math.*
- Ohlson, M., Z. Andrushchenko, and D. von Rosen (2008). Explicit estimators of the parameters in a multivariate normal distribution when the covariance matrix is banded of order m . *Research Report 2008:3, Centre of Biostochastics. Swedish University of Agricultural Sciences.*
- von Rosen, D. (1995). Residuals in the growth curve model. *Ann. Inst. Statist. Math.* 47, 129–136